

NON-FACTORIAL NODAL COMPLETE INTERSECTION THREEFOLDS

SLAWOMIR CYNK AND SLAWOMIR RAMS

ABSTRACT. We give a bound on the minimal number of singularities of a nodal projective complete intersection threefold which contains a smooth complete intersection surface that is not a Cartier divisor.

INTRODUCTION

The main subject of this consideration are non-factorial nodal complete intersections in projective space. Our interest in nodal varieties is justified by the fact that an ordinary double point (a node) is the most ubiquitous singularity that appears in analytic, algebraic geometry and singularity theory. A remarkable property of three-dimensional nodes is the existence of the so-called small resolution, i.e. an analytic modification that replaces the singular point with one copy of projective line. In general a small resolution exists only as an analytic space. Certain global properties of a nodal variety (e.g. existence of a Kähler small resolution, topological invariants) are closely related to the structure of its class group of Weil divisors. Locally analytically for every node there exists germ of a smooth surface on the threefold that passes through the singularity. Is a subtle question whether algebraic surfaces with such a property exist on a given threefold.

A variety is called factorial if every Weil divisor on it is Cartier. In the case of a complete intersection threefold X in projective space it means that each surface on X is defined by a single homogeneous polynomial. Cheltsov ([3], [4]), Cheltsov and Park ([5]), Kosta ([12], [13]) gave some lower bounds on the number of singular points on a non-factorial threefold hypersurfaces in $\mathbb{P}_4$ and complete intersections in $\mathbb{P}_5$ in terms of degrees of defining equations.

For a smooth surface in projective space a generic threefold complete intersection of hypersurfaces of sufficiently high degrees that contain the surface is nodal (see e.g. [11]). Our goal is to study this construction in more detail in the case of a complete intersection surface (Thm. 3.1). The number of nodes of the considered threefold can be computed using Chern classes (see Prop. 2.4). Using the above fact and an elementary but quite tedious integer inequality (Lemma 1.1) we obtain our main result.

Theorem 2.1 *Let $X \subset \mathbb{P}_{k+3}(\mathbb{C})$ be a nodal complete intersection threefold and let $I(X) = \langle f_1, \dots, f_k \rangle$, where the sequence $(\deg(f_1), \dots, \deg(f_k))$ is non-decreasing. Assume that*

the set $V(f_1, \dots, f_s)$ is smooth in codimension 3 for $s \leq (k-1)$,

If a smooth complete intersection surface is not a Cartier divisor on X , then either

$$\# \text{sing}(X) \geq \sum_{1 \leq i \leq j \leq k} (\deg(f_i) - 1) \cdot (\deg(f_j) - 1)$$

or

$$\# \text{sing}(X) \geq 2^{k-1} \text{ and } k \in \{2, 3, 4\}, \deg(f_1) = \dots = \deg(f_k) = 2.$$

2010 *Mathematics Subject Classification.* Primary: 32S20; Secondary 14M10.
Research partially supported by NCN grant no. N N201 608040.

Motivated by the above result we formulate Conjecture (see p. 6), which in the case of a complete intersection in $\mathbb{P}_5$ is a slightly stronger version of [2, Conjecture 30] (see Remark 2.7).

In the last section of the paper we construct numerous examples which show that all integer sequences considered in Sect. 2 can be realized geometrically with help of complete intersections (Thm. 3.1).

1. AN INTEGER INEQUALITY

Given two non-decreasing sequences of positive integers $\underline{d} := (d_1, \dots, d_k)$ and $\underline{e} := (e_1, \dots, e_{k+1})$ we define the following quantity

$$(1) \quad \mathcal{S}(\underline{d}; \underline{e}) := e_1 \cdot \dots \cdot e_{k+1} \cdot \sum_{1 \leq i \leq j \leq k} (d_i - e_i) \cdot (d_j - e_{j+1}).$$

The goal of this section is to prove an elementary inequality involving $\mathcal{S}(\cdot; \cdot)$ that will be useful in the sequel. We work under the additional assumption that $(\underline{d}; \underline{e})$ satisfies the following conditions:

- (2) $d_i \geq e_i \quad \text{for } i \leq k,$
- (3) $d_i < e_{i+1} \Rightarrow (d_1, \dots, d_i) = (e_1, \dots, e_i), \text{ for } 1 \leq i \leq k,$
- (4) $d_i < e_{i+2} \Rightarrow (d_1, \dots, d_i) \subset (e_1, \dots, e_{i+1}) \text{ for } 1 \leq i \leq k-1,$

where, by abuse of notation, we write $(d_1, \dots, d_i) \subset (e_1, \dots, e_{i+1})$ iff $(d_1, \dots, d_i)$ is a subsequence of $(e_1, \dots, e_{i+1})$. In particular, the conditions (2), (3) imply that each summand in $\mathcal{S}(\underline{d}; \underline{e})$ is non-negative. Moreover, by direct computation one obtains

$$(5) \quad \mathcal{S}(\underline{d}; \underline{e}) = 0 \quad \text{iff} \quad \underline{d} \subset \underline{e}.$$

Furthermore, we have the following elementary observation

$$(6) \quad \text{if } d_i > e_{i+1} \text{ for } i = 1, \dots, k \quad \text{then} \quad \mathcal{S}(\underline{d}; \underline{e}) \geq \mathcal{S}(\underline{d}; 1^{k+1}),$$

where, to simplify our notation, for an integer $c \in \mathbb{N}$ we put $c^k := \underbrace{c, \dots, c}_{k \text{ times}}$. Indeed, the inequalities

$1 \leq e_i \leq e_{i+1} < d_i$ imply that

$$(7) \quad e_i \cdot (d_i - e_i) \geq (d_i - 1) \quad \text{and} \quad e_{i+1} \cdot (d_i - e_{i+1}) \geq (d_i - 1).$$

Now (6) follows from (7) and (1).

After those preparations we can prove the main result of this section.

Lemma 1.1. *Let $\underline{d} = (d_1, \dots, d_k)$, $\underline{e} = (e_1, \dots, e_{k+1})$ be non-decreasing sequences of positive integers such that $d_1 \geq 2$ and the conditions (2), (3), (4) are fulfilled. Then either*

$$\mathcal{S}(\underline{d}; \underline{e}) = 0$$

or

$$(8) \quad \mathcal{S}(\underline{d}; \underline{e}) \geq \mathcal{S}(\underline{d}; 1^{k+1})$$

or

$$\mathcal{S}(\underline{d}; \underline{e}) = 2^{k-1} \quad \text{and } k \in \{2, 3, 4\}, \quad e_1 = e_2 = 1, \quad d_1 = \dots = d_k = e_3 = \dots = e_{k+1} = 2.$$

Proof. We will proceed by induction on k , the case $k = 1$ follows from the observations (5) and (6). Now, assume that the lemma holds for $k - 1$ and fix sequences $\underline{d}$ and $\underline{e}$ satisfying the assumptions of the lemma. We shall separately consider the case $d_1 = e_1$. If $(d_2, \dots, d_k) \subset (e_2, \dots, e_{k+1})$,

then $\mathcal{S}(\underline{d}; \underline{e}) = 0$ by (5). Otherwise, as the sequences $(d_2, \dots, d_k)$ and $(e_2, \dots, d_{k+1})$ satisfy the assumptions of the lemma, we can apply the induction hypothesis (8) and obtain

$$\mathcal{S}(\underline{d}; \underline{e}) = d_1 \cdot \mathcal{S}(d_2, \dots, d_k; e_2, \dots, e_{k+1}) \geq d_1 \cdot \mathcal{S}(d_2, \dots, d_k; \underline{1}_k) = \mathcal{S}(\underline{d}; \underline{1}^k, d_1).$$

Since the equality $d_k = d_1$ would imply $(\underline{d} \subset \underline{e})$, we have $d_k > d_1$ and (6) gives

$$\mathcal{S}(\underline{d}; \underline{1}^k, d_1) \geq \mathcal{S}(\underline{d}; \underline{1}^{k+1}).$$

This completes the proof in the case $d_1 = e_1$.

It remains to consider the case $d_1 > e_1$. The condition (3) yields $d_i \geq e_{i+1}$ for each $i \leq k$, and using (7) we obtain

$$(9) \quad \mathcal{S}(\underline{d}; \underline{e}) \geq \mathcal{S}(\underline{d}; 1, e_2, \dots, e_{k+1}).$$

By (1) we have

$$(10) \quad \mathcal{S}(\underline{d}; 1, e_2, \dots, e_{k+1}) = e_2 \cdot \dots \cdot e_{k+1} \cdot (d_1 - 1) \cdot \sum_{j=1}^k (d_j - e_{j+1}) + \mathcal{S}(d_2, \dots, d_k; e_2, \dots, e_{k+1}).$$

If $\sum_{j=1}^k (d_j - e_{j+1}) \leq 0$, then the inequalities $d_i \geq e_{i+1}$, where $i = 1, \dots, k$, yield $\underline{d} \subset \underline{e}$. Otherwise, repeated use of (7) gives

$$(11) \quad e_2 \cdot \dots \cdot e_{k+1} \cdot (d_1 - 1) \cdot \sum_{j=1}^k (d_j - e_{j+1}) \geq (d_1 - 1) \cdot \left(\sum_{j=1}^k d_j - k \right).$$

Now, we are in position to apply the induction hypothesis.

Suppose that $\mathcal{S}(d_2, \dots, d_k; e_2, \dots, e_{k+1}) \geq \mathcal{S}(d_2, \dots, d_k; \underline{1}^k)$. Then (9), (10) and (11) yield

$$\mathcal{S}(\underline{d}; \underline{e}) \geq (d_1 - 1) \cdot \left(\sum_{j=1}^k d_j - k \right) + \mathcal{S}(d_2, \dots, d_k; \underline{1}^k) = \mathcal{S}(\underline{d}; \underline{1}^{k+1}).$$

Assume, that $\mathcal{S}(d_2, \dots, d_k; e_2, \dots, e_{k+1})$ vanishes. Then, by (5), we have

$$(d_2, \dots, d_k) \subset (e_2, \dots, e_{k+1}).$$

Therefore, for $i = 2, \dots, k$ we have either $d_i = e_i$ or $d_i = e_{i+1}$. Since $\underline{e}$ is non-decreasing, we obtain

$$d_i \leq e_{i+1} \text{ for } i = 2, \dots, k.$$

On the other hand, the assumption $d_1 > e_1$ combined with (3) gives $d_i \geq e_{i+1}$ for $i = 1, \dots, k$. Thus we have $d_i = e_{i+1}$ for $i = 2, \dots, k$.

If $d_{k-1} < d_k = e_{k+1}$, then the condition (4) gives $(d_1, \dots, d_{k-1}) \subset (e_1, e_2, d_2, \dots, d_{k-1})$. Since $d_1 > e_1$, we have $d_1 = e_2$ and $\underline{d} \subset \underline{e}$. Consequently, by (5), we can restrict our attention to the case $d_{k-1} = d_k$. Repeating this reasoning we obtain that either $\mathcal{S}(\underline{d}; \underline{e}) = 0$ or

$$d_1 = \dots = d_k = e_3 = \dots = e_{k+1} \text{ and } e_2 < d_1.$$

Suppose that $d_1 \geq 3$. Using (7) we obtain the inequalities

$$\mathcal{S}(d_1, \dots, d_1; e_1, e_2, d_1, \dots, d_1) \geq d_1^{k-1} \cdot (d_1 - 1)^2 \geq \binom{k+1}{2} \cdot (d_1 - 1)^2 = \mathcal{S}(d_1, \dots, d_1; \underline{1}^{k+1}).$$

For $d_1 = 2$ we get $e_1 = e_2 = 1$, so

$$\mathcal{S}(2^k; 1, 1, 2^{k-1}) = 2^{k-1} \geq \binom{k+1}{2} = \mathcal{S}(2^k; \underline{1}^{k+1}) \text{ for } k \geq 5.$$

Finally, assume that

$$(d_2, \dots, d_k; e_2, \dots, e_{k+1}) \in \{(2^2; 1^2, 2), (2^3; 1^2, 2^2), (2^4; 1^2, 2^3)\}$$

is one of the exceptional cases. Then $d_1 = 2$, $e_1 = 1$, so $(\underline{d}, \underline{e}) \in \{(2^3; 1^3, 2), (2^4; 1^3, 2^2), (2^5; 1^3, 2^3)\}$. By direct computation (8) holds in these cases. $\square$

2. MAIN RESULT

In this section we prove the main result of this note

Theorem 2.1. *Let $X \subset \mathbb{P}_{k+3}(\mathbb{C})$ be a nodal complete intersection threefold and let $I(X) = \langle f_1, \dots, f_k \rangle$, where the sequence $(\deg(f_1), \dots, \deg(f_k))$ is non-decreasing. Assume that*

$$(12) \quad \text{the set } V(f_1, \dots, f_s) \text{ is smooth in codimension 3 for } s \leq (k-1),$$

If a smooth complete intersection surface is not a Cartier divisor on X , then either

$$\# \text{sing}(X) \geq \sum_{1 \leq i \leq j \leq k} (\deg(f_i) - 1) \cdot (\deg(f_j) - 1)$$

or

$$\# \text{sing}(X) \geq 2^{k-1} \text{ and } k \in \{2, 3, 4\}, \deg(f_1) = \dots = \deg(f_k) = 2.$$

Remark 2.2. The local class group of a node is $\mathbb{Z}$ (see [15]), so the nodal threefold is factorial iff it is $\mathbb{Q}$ -factorial. Consequently, Thm 2.1 gives a bound for non- $\mathbb{Q}$ -factorial complete intersections. For the sake of simplicity in this paper we formulate all results using the notion of factoriality.

Remark 2.3. Obviously the equations defining a complete intersection are not unique, whereas their degrees are uniquely determined. The latter can be computed f.i. from the minimal resolution.

In order to simplify our notation, for a nodal complete intersection threefold $X \subset \mathbb{P}_{k+3}(\mathbb{C})$ and a smooth complete intersection surface $S \subset X$ from now on we put

$$(13) \quad I(X) = \langle f_1, \dots, f_k \rangle \quad \text{with} \quad \deg(f_1) \leq \dots \leq \deg(f_k),$$

$$(14) \quad I(S) = \langle g_1, \dots, g_{k+1} \rangle \quad \text{with} \quad \deg(g_1) \leq \dots \leq \deg(g_{k+1}).$$

The proof of Thm 2.1 will be preceded by a proposition and a lemma.

Proposition 2.4. (cf. [1, Ex. 5]) *Let $X \subset \mathbb{P}_{k+3}(\mathbb{C})$ be a nodal complete intersection threefold and let $S \subset X$ be a smooth complete intersection surface. Then X has exactly*

$$\mathcal{S}(\deg(f_1), \dots, \deg(f_k); \deg(g_1), \dots, \deg(g_{k+1}))$$

nodes on S .

Proof. Let $X := Y_1 \cap \dots \cap Y_k$, where $Y_i = V(f_i)$. Let $\sigma : \tilde{\mathbb{P}}_{k+3} \rightarrow \mathbb{P}_{k+3}$ be the blow-up of $\mathbb{P}_{k+3}$ along the smooth surface S and let $\tilde{X}$ be the strict transform of X . Then $\tilde{X}$ is again a complete intersection, i.e. $\tilde{X} = \tilde{Y}_1 \cap \dots \cap \tilde{Y}_k$, where $\tilde{Y}_i$ is the strict transform of Y_i . Since $\tilde{X}$ is a small resolution of the nodes of X that lie on S , we have

$$\tilde{e}(\tilde{X}) = e(X) + \nu,$$

where $\nu := \#(\text{sing}(X) \cap S)$ and e (resp. $\tilde{e}$) denotes the topological Euler characteristic (resp. degree of the Fulton-Johnson class) as in [1]. On the other hand we have

$$e(X) = \tilde{e}(X) + \# \text{sing}(X),$$

$$e(\tilde{X}) = \tilde{e}(\tilde{X}) + \# \text{sing}(X) - \nu,$$

because the Milnor number of a node is one. Consequently

$$\nu = \frac{1}{2}(\tilde{e}(\tilde{X}) - \tilde{e}(X)).$$

By [1, Prop. 1] the number $(\tilde{e}(\tilde{X}) - \tilde{e}(X))$ is a polynomial in $\deg(f_1), \dots, \deg(f_k), \deg(g_1), \dots, \deg(g_{k+1})$ and equals

$$\mathcal{S}(\deg(f_1), \dots, \deg(f_k); \deg(g_1), \dots, \deg(g_{k+1}))$$

if $\deg(f_i) \geq \deg(g_j)$ for all $1 \leq i \leq k$ and $1 \leq j \leq (k+1)$. Consequently, the claim of the lemma holds. $\square$

In the next lemma we maintain the notation (13), (14).

Lemma 2.5. *Let $X \subset \mathbb{P}_{k+3}(\mathbb{C})$ be a nodal complete intersection threefold, such that (12) holds, and let $S \subset X$ be a smooth complete intersection surface.*

(I) *For $j \leq k$ one has the inequality:*

$$\deg(f_j) \geq \deg(g_j).$$

(II) *If $\deg(f_i) < \deg(g_{i+1})$ for an integer $i \leq k$, then*

$$(\deg(f_1), \dots, \deg(f_i)) = (\deg(g_1), \dots, \deg(g_i)).$$

(III) *If $\deg(f_i) < \deg(g_{i+2})$ for an integer $i \leq (k-1)$, then*

$$(\deg(f_1), \dots, \deg(f_i)) \subset (\deg(g_1), \dots, \deg(g_{i+1})).$$

Proof. (I) Let i_0 be the smallest integer such that $\deg(f_{i_0}) < \deg(g_{i_0})$. Observe that $i_0 > 1$. Indeed, otherwise $\deg(f_1) < \deg(g_l)$ for $l = 1, \dots, k+1$, which implies $f_1 \notin I(S)$ and contradicts the assumption that S is contained in X .

Since $I(X) \subset I(S)$, we have $f_j \in \langle g_1, \dots, g_{k+1} \rangle$ for each j . On the other hand, if $i \leq i_0$ then the inequality

$$\deg(f_i) \leq \deg(f_{i_0}) < \deg(g_{i_0})$$

holds, which implies $f_j \in \langle g_1, \dots, g_{i_0-1} \rangle$. We obtain the inclusion

$$V(g_1, \dots, g_{i_0-1}) \subset V(f_1, \dots, f_{i_0}).$$

The latter contradicts the assumption that X is a complete intersection threefold and completes the proof of part (I).

(II) Assume that $\deg(f_i) < \deg(g_{i+1})$. We repeat the proof of part (I) to obtain the inclusion

$$\langle f_1, \dots, f_i \rangle \subset \langle g_1, \dots, g_i \rangle.$$

Consequently, there exist homogeneous polynomials $h_{l,j}$ such that

$$(15) \quad f_l = \sum_{j=1}^i h_{l,j} g_j \quad \text{for } l = 1, \dots, i.$$

Obviously, one has $h_{l,j} = 0$ when $\deg(f_l) < \deg(g_j)$. Otherwise, we have

$$(16) \quad \deg(h_{l,j}) = (\deg(f_l) - \deg(g_j)).$$

Fix a point $P \in V(g_1, \dots, g_i)$. Since the surface S is assumed to be smooth, the polynomials $g_1, \dots, g_i$ form part of a coordinate system around P on $\mathbb{P}_{k+3}$. Observe, that the Jacobi matrix of $f_1, \dots, f_i$ with respect to such coordinates in the point P becomes

$$\begin{bmatrix} h_{1,1} & \dots & h_{1,i} & 0 & \dots & 0 \\ \vdots & & \vdots & \vdots & & \vdots \\ h_{i,1} & \dots & h_{i,i} & 0 & \dots & 0 \end{bmatrix}.$$

In particular, if we put $H := [h_{l,j}]_{l=1, \dots, i}^{j=1, \dots, i}$, then we have

$$(17) \quad V(g_1, \dots, g_i, \det(H)) \subset \text{sing}(V(f_1, \dots, f_i)).$$

Suppose that $\deg(\det(H)) > 0$. Then, $\text{sing}(V(f_1, \dots, f_i))$ has a codimension-one component, which is impossible by (12). Therefore, (16) yields

$$\sum_{j=1}^i (\deg(f_j) - \deg(g_j)) = \deg(\det(H)) = 0.$$

The claim results from part (I) of the lemma.

(III) Assume that $\deg(f_i) < \deg(g_{i+2})$. As in the proof of part (I) we obtain the inclusion

$$\langle f_1, \dots, f_i \rangle \subset \langle g_1, \dots, g_{i+1} \rangle$$

and the homogeneous polynomials $h_{l,j}$ such that

$$(18) \quad f_l = \sum_{j=1}^{i+1} h_{l,j} g_j \quad \text{for } l = 1, \dots, i.$$

Moreover, either (16) holds or $h_{l,j} = 0$ when $\deg(f_l) < \deg(g_j)$.

Put $Z := \{x \in \mathbb{P}_{k+3} : g_1(x) = \dots = g_{i+1}(x) = 0 \text{ and } \text{rank}(H(x)) \leq (i-1)\}$, where $H := [h_{l,j}]_{l=1, \dots, i}^{j=1, \dots, i+1}$. As in (17) we have the inclusion

$$Z \subset \text{sing}(V(f_1, \dots, f_i)).$$

By [8, Prop. 17.25] either $Z = \emptyset$ or $\dim(Z) = (k-i)$. Thus the assumption (12) yields $Z = \emptyset$. In particular, the set $V(f_1, \dots, f_i)$ is smooth along $V(g_1, \dots, g_{i+1})$, so the latter is a Cartier divisor on $V(f_1, \dots, f_i)$. By [9, Cor. IV.3.2] $V(g_1, \dots, g_{i+1})$ is cut out by a hyperplane from $V(f_1, \dots, f_i)$, which completes the proof of (III). $\square$

After those preparations we can give the proof of Thm 2.1:

Proof of Thm 2.1. We maintain the notation (13), (14). Observe that without loss of generality we can assume that X is non-degenerate.

Since $S \subset X$ is not Cartier, the threefold X has a node on S . By Prop. 2.4 we have

$$\mathcal{S}(f_1, \dots, f_k; g_1, \dots, g_{k+1}) > 0.$$

Lemma 2.5 yields that the non-decreasing sequences $(f_1, \dots, f_k)$, $(g_1, \dots, g_{k+1})$ satisfy the conditions (2), (3), (4). Finally, the theorem results directly from Lemma 1.1. $\square$

Motivated by Thm. 2.1 we propose the following conjecture:

Conjecture. *Let $X := V(f_1, \dots, f_k) \subset \mathbb{P}_{k+3}$ be a nodal complete intersection threefold such that the sequence $(\deg(f_1), \dots, \deg(f_k))$ is non-decreasing and $V(f_1, \dots, f_i)$ is smooth in codimension three for $i = 1, \dots, k-1$. If X is not factorial, then either*

$$\# \text{sing}(X) \geq \sum_{1 \leq i \leq j \leq k} (\deg(f_i) - 1) \cdot (\deg(f_j) - 1)$$

or

$$\# \text{sing}(X) \geq 2^{k-1} \text{ and } k \in \{2, 3, 4\}, \deg(f_1) = \dots = \deg(f_k) = 2.$$

We shall see in the next section that if the above conjecture holds true then the bound it provides is sharp (see Thm. 3.6). For the hypersurface case the conjecture was proved by I. Cheltsov ([4]). For a complete intersection in $\mathbb{P}_5$ Cheltsov formulated similar conjecture (see [2, Conj. 30]). He puts a stronger assumption on X , namely he assumes that the hypersurface $V(f_1)$ is smooth, whereas Thm. 2.1 provides a strong evidence that the bound holds also if $V(f_1)$ is allowed to have isolated singularities. The necessity of an assumption on the singularities of $V(f_1)$ has been observed by Cheltsov in the following example.

Example 2.6 ([2, Ex. 29]). Let $Y_1 = V(l_1 l_3 + l_2 l_4)$ be a rank-4 quadric in $\mathbb{P}_5$, and let $Y_2 = V(f_2)$ be a general degree- n hypersurface. Then the complete intersection $X := Y_1 \cap Y_2$ has exactly n nodes in $V(l_1, l_2, l_3, l_4, f_2)$ as its only singularities. The threefold X is not factorial (as it contains the smooth surface $V(l_1, l_3, f_2)$), while

$$\# \text{sing}(X) = n < n^2 - n + 1,$$

which violates the bound of Thm. 2.1.

Observe that Y_1 is singular along the line $L := V(l_1, \dots, l_4)$, and each singularity of X is obtained as a point where L and Y_2 meet. The above complete intersection corresponds to the sequences $\underline{d} = (2, n)$, $\underline{e} = (1, 1, n)$ that do not satisfy the condition (4).

For a fixed integer $k > 2$ and $n \gg k$ the sequences $\underline{d} := (2^{k-1}, n)$ and $\underline{e} := (1^k, n)$ produce (via Prop. 3.6) similar examples in $\mathbb{P}_{k+3}$.

Remark 2.7. Observe that Thm. 2.1 implies that the above Cheltsov's example with $n = 2$ and f_1, f_2 interchanged (i.e. $V(f_2)$ considered as the first hypersurface) is the only counterexample to [2, Conjecture 30] in which a complete intersection surface fails to be a Cartier divisor. The example in question is one of the three exceptional cases in Thm. 2.1 (i.e. $k = 2$, $e_1 = e_2 = 1$ and $e_3 = d_1 = d_2 = 2$).

3. GENERAL COMPLETE INTERSECTIONS

In this section we prove the existence of nodal complete intersection threefolds that appear in Thm. 2.1. Namely, we show the following theorem.

Theorem 3.1. *Let $\underline{d}, \underline{e}$ be a pair of non-decreasing sequences of positive integers satisfying the conditions (2)–(4). Then there exists a smooth surface $S = V(g_1, \dots, g_{k+1})$ and a threefold $X = V(f_1, \dots, f_k)$ with $\deg(f_i) = d_i$, $\deg(g_j) = e_j$ such that $S \subset X$ and the threefold X has exactly $\mathcal{S}(\underline{d}; \underline{e})$ nodes on S as its only singularities.*

In particular, if $\mathcal{S}(\underline{d}, \underline{e}) > 0$, then the surface S is not a Cartier divisor, hence the variety X is not factorial.

In fact we shall show a more explicit statement (Prop. 3.6). In order to avoid unnecessary technical complications in its proof, we represent the polynomial f_i as a linear combination of minimal number of the generators g_j . To make this explicit, for $i \leq (k-1)$ we define

$$\alpha_{(\underline{d}; \underline{e})}(i) := \begin{cases} i & \text{if } d_i < e_{i+1}, \\ i+1 & \text{if } d_i \geq e_{i+1} \text{ and } d_i < e_{i+2}, \\ i+2 & \text{otherwise.} \end{cases}$$

Whenever it leads to no ambiguity we write $\alpha(i) := \alpha_{(\underline{d}; \underline{e})}(i)$. Finally, we put $\alpha(k) := k+1$.

We choose non-zero homogeneous polynomials $g_1, \dots, g_{k+1}$ such that $\deg(g_j) = e_j$ and

$$(19) \quad V(g_1, \dots, g_i) \text{ is smooth for } i \leq k+1$$

Given the polynomials $g_1, \dots, g_{k+1}$ and homogeneous polynomials $h_{i,j}$ of degree $(d_i - e_j)$, we define a degree- d_i homogeneous polynomial $f_i \in \langle g_1, \dots, g_{\alpha(i)} \rangle$ by the equality

$$(20) \quad f_i = \sum_{j=1}^{\alpha(i)} h_{i,j} g_j.$$

In order to prove Prop. 3.6 we need several lemmata. In their proofs we put $h_{i,j} := 0$ for $j > \alpha(i)$ and define the matrix

$$H_i := [h_{l,j}]_{l=1, \dots, i}^{j=1, \dots, \alpha(i)}.$$

At first we deal with the smooth case.

Lemma 3.2. Fix $g_1, \dots, g_{k+1}$ that satisfy (19). If $i \leq k-1$ and $\alpha(i) \leq i+1$, then for generic polynomials $h_{l,j}$, where $l \leq i$ and $j \leq \alpha(l)$, the intersection $V(f_1, \dots, f_i)$ is smooth along $V(g_1, \dots, g_{\alpha(i)})$.

Proof. Case $\alpha(i) = i$: As in the proof of Lemma 2.5.(II) we have the equality

$$V(g_1, \dots, g_i, \det(H_i)) = \text{sing}(V(f_1, \dots, f_i)) \cap V(g_1, \dots, g_i)$$

and $\det(H_i)$ is a degree-0 polynomial. Obviously, for generic $h_{l,j}$, we have $\det(H_i) \neq 0$.

Case $\alpha(i) = i+1$: As in the proof of Lemma 2.5 we have the equality

$$(21) \quad \text{sing}(V(f_1, \dots, f_i)) \cap V(g_1, \dots, g_{i+1}) = \{P \in V(g_1, \dots, g_{i+1}) : \text{rank}(H_i(P)) \leq (i-1)\}.$$

Moreover, by (4), there exists an integer $i_0 \geq 1$ such that

$$(e_1, \dots, e_{k+1}) = (d_1, \dots, d_{i_0}, e_{i_0+1}, d_{i_0+1}, \dots, d_k),$$

so the polynomials $h_{1,1}, \dots, h_{i_0, i_0}, h_{i_0+1, i_0+2}, \dots, h_{i, i+1}$ are constant and determinant of the matrix obtained from H_i by deleting its i_0^{th} column is a degree-0 polynomial. For generic $h_{l,j}$ the determinant in question does not vanish and the lemma follows. $\square$

Lemma 3.3. Assume that $i \leq (k-1)$ and $\alpha(i) \leq (i+1)$. For generic choice of the polynomials $g_1, \dots, g_{\alpha(i)}$ and $h_{l,j}$, where $l \leq i$ and $j \leq \alpha(l)$ the intersection $V(f_1, \dots, f_i)$ is smooth.

Proof. We proceed by induction on i .

[i=1]: If $\alpha(1) = 1$, then $f_1 = h_{1,1} \cdot g_1$ with $\deg(h_{1,1}) = 0$, so $V(g_1) = V(f_1)$ is smooth by assumption. Suppose that $\alpha(1) = 2$, then $f_1 = h_{1,1} \cdot g_1 + h_{1,2} \cdot g_2$ with $\deg(h_{1,2}) = 0$. Fix g_1, g_2 satisfying (19). By Bertini, for generic $h_{1,1}, h_{1,2}$ the hypersurface $V(f_1)$ is smooth away from $V(g_1, g_2)$. Therefore, generic $V(f_1)$ is smooth by Lemma 3.2.

[(i-1) $\rightsquigarrow$ i]: If $\alpha(i-1) \leq i$, then by the induction hypothesis, for generic $g_1, \dots, g_i$ and $h_{1,1}, \dots, h_{i-1, i}$ the intersection $V(f_1, \dots, f_{i-1})$ is smooth. By Bertini, for generic $h_{i,1}, \dots, h_{i,i}$ the intersection $V(f_1, \dots, f_i)$ is smooth away from $V(g_1, \dots, g_i)$. Lemma 3.2 completes the proof in this case.

Suppose that $\alpha(i-1) = i+1$. By definition, we have $d_{i-1} \geq e_{i+1}$. From $\alpha(i) = i+1$ and (4) we infer that $d_i \in \{e_i, e_{i+1}\}$, so $d_i \leq e_{i+1}$ and $d_{i-1} = d_i = e_{i+1}$. Finally, from (4) we obtain $d_{i-1} \in \{e_{i-1}, e_i\}$, so $d_{i-1} \leq e_i \leq e_{i+1}$ and $d_{i-1} = e_i$. In particular, we have the inclusion

$$(d_1, \dots, d_{i-1}) \subset (e_1, \dots, e_i).$$

Consequently, if we define $\tilde{d} := (d_1, \dots, d_{i-1}, d_{i-1}+1, \dots, d_{i-1}+k-i+1)$ and $\tilde{e} := (e_1, \dots, e_i, d_{i-1}+1, \dots, d_{i-1}+k-i+1)$, then the pair $(\tilde{d}, \tilde{e})$ satisfies the conditions (2), (3), (4). By definition, for $l < i-1$

$$\alpha_{(\tilde{d}, \tilde{e})}(l) = \alpha_{(\underline{d}, \underline{e})}(l) \text{ and } \alpha_{(\tilde{d}, \tilde{e})}(i-1) = i$$

so, by the induction hypothesis, $V(\tilde{f}_1, \dots, \tilde{f}_{i-1})$ is smooth if we make generic choices of $\tilde{h}_{l,j}$, where $l \leq i-1$ and $j \leq \alpha_{(\tilde{d}, \tilde{e})}(l)$. Assume the latter holds, and put $h_{l,j} = \tilde{h}_{l,j}$ for $l \leq i-1$ and $j \leq \alpha_{(\tilde{d}, \tilde{e})}(l)$. Then for generic choice of $h_{i-1, i+1} \in \mathbb{C}$ the intersection $V(f_1, \dots, f_{i-1})$ is smooth. Finally, Bertini and Lemma 3.2 imply that $V(f_1, \dots, f_i)$ is smooth for generic choices of $h_{i,1}, \dots, h_{i, i+1}$. $\square$

Now we can deal with the general case.

Lemma 3.4. Let $i \leq (k-1)$. For generic choice of the polynomials $g_1, \dots, g_{\alpha(i)}$ and $h_{l,j}$, where $l \leq i$ and $j \leq \alpha(l)$, the intersection $V(f_1, \dots, f_i)$ is smooth in codimension four.

Proof. We proceed by induction on i .

[i=1]: If $\alpha(1) \leq 2$, then generic $V(f_1)$ is smooth by Lemma 3.3. Suppose that $\alpha(1) = 3$, then

$$f_1 = h_{1,1} \cdot g_1 + h_{1,2} \cdot g_2 + h_{1,3} \cdot g_3.$$

If (19) holds, we have $\text{sing}(V(f_1)) \subset V(g_1, g_2, g_3, h_{1,1}, h_{1,2}, h_{1,3})$ for generic $h_{1,1}, \dots, h_{1,3}$.

[(i-1) $\rightsquigarrow$ i]: By Lemma 3.3 we can assume $\alpha(i) = i + 2$. We apply Bertini on $\text{reg}(V(f_1, \dots, f_{i-1}))$ to obtain the inclusion

$$\text{sing}(V(f_1, \dots, f_i)) \subset V(g_1, \dots, g_{i+2}) \cup \text{sing}(V(f_1, \dots, f_{i-1})).$$

For generic choice of g_{i+2} no component of $\text{sing}(V(f_1, \dots, f_{i-1}))$ is contained in $V(g_{i+2})$. By the induction hypothesis, it remains to study the components of $\text{sing}(V(f_1, \dots, f_i))$ that are contained in $V(g_1, \dots, g_{i+2})$.

Recall that we assumed (19), so we have the equality

$$\text{sing}(V(f_1, \dots, f_i)) \cap V(g_1, \dots, g_{i+2}) = \{P \in V(g_1, \dots, g_{i+2}) : \text{rank}(H_i(P)) \leq (i-1)\}.$$

In particular, the induction hypothesis implies that for generic g_{i+2} the set

$$(22) \quad \{P \in V(g_1, \dots, g_{\alpha(i-1)}) : \text{rank}(H_{i-1}(P)) \leq (i-2)\}$$

has codimension at least five in $V(f_1, \dots, f_{i-1}, f_i)$. Thus, we can restrict our attention to the components of the singular locus that meet the set $\{P : \text{rank}(H_{i-1}(P)) = (i-1)\}$.

Observe that $\alpha(i-1) < i+2$, so the components in question are contained in $V(h_{i,i+2})$. If $\deg(h_{i,i+2}) = 0$ we can choose a nonzero $h_{i,i+2}$ and the proof is complete. Otherwise, we can assume that the hypersurface $V(h_{i,i+2})$ meets $V(g_1, \dots, g_{i+2})$ properly.

Let $\hat{H}_{i-1} := [h_{l,j}]_{l=1, \dots, i-1}^{j=1, \dots, i-1}$. Determinant of the minor of H_i obtained by removing the i^{th} and $(i+2)^{\text{nd}}$ column can be written as

$$(23) \quad h_{i,i+1} \cdot \det(\hat{H}_{i-1}) + h_{i-1,i+1} \cdot \det(M_1),$$

where M_1 is another minor of H_i . One can easily see, that for generic $h_{i,1}, \dots, h_{i,i-1}$ and $h_{i,i+1}$ the hypersurface given by (23) meets $V(g_1, \dots, g_{i+2}, h_{i,i+2})$ properly. An analogous argument applied to the minor obtained by removing the $(i+1)^{\text{st}}$ and $(i+2)^{\text{nd}}$ column from H_i shows that the

$$\text{codim}(\text{sing}(V(f_1, \dots, f_i)) \cap \{P : \text{rank}(H_{i-1}(P)) = (i-1)\}) \geq 5,$$

and it remains to deal with the components of $\text{sing}(V(f_1, \dots, f_i))$ that are contained in the set

$$(24) \quad V(g_1, \dots, g_{i+2}, h_{i,i+2}, \det(\hat{H}_{i-1})).$$

If $\alpha(i-1) = i-1$, then $\det(\hat{H}_{i-1})$ is constant and we are done. Otherwise, we can assume that (24) is of codimension four and use another minor of the matrix H_{i-1} to complete the proof. $\square$

Lemma 3.5. *Assume that $g_1, \dots, g_{k+1}$ satisfy the condition (19) and the fourfold $V(f_1, \dots, f_{k-1})$ is smooth. For generic $f_k \in \mathcal{O}_{\mathbb{P}_{k+3}}(d_k) \otimes \mathcal{I}(S)$ the intersection $V(f_1, \dots, f_k)$ has at most nodes (A_1 singularities).*

Proof. Put $Y := V(f_1, \dots, f_{k-1})$. Bertini theorem implies that generic $V(f_k)$ is smooth away from the surface S . We cover the surface in question with the open sets

$$U_{i,j} := \{P \in S : \mathcal{I}(S)\mathcal{O}_{Y,P} = \langle g_i, g_j \rangle \mathcal{O}_{Y,P}\},$$

where $i, j = 1, \dots, k+1$. By stability property of A_1 points, the set

$$\mathcal{L}_{i,j} := \{f_k \in \mathcal{O}_{\mathbb{P}_{k+3}}(d_k) \otimes \mathcal{I}(S) : V(f_k|_Y) \text{ has at most nodes on } U_{i,j}\}$$

is open in $\mathcal{O}_{\mathbb{P}_{k+3}}(d_k) \otimes \mathcal{I}(S)$. To show that $\mathcal{L}_{i,j}$ is non-empty observe that for generic degree- $(d_k - e_i)$ (resp. degree- $(d_k - e_j)$) homogeneous polynomial $h_{k,i}$ (resp. $h_{k,j}$) the threefold given on Y by vanishing of

$$(25) \quad h_{k,i} \cdot g_i + h_{k,j} \cdot g_j$$

has nodes in $V(g_i, g_j, h_{k,i}, h_{k,j})$ as its only singularities. In particular, the polynomial (25) belongs to $\mathcal{L}_{i,j}$. $\square$

Now we are in position to formulate and prove the following proposition that directly implies Thm. 3.1. In Prop. 3.6 we maintain the notation (20).

Proposition 3.6. *Let $\underline{d} := (d_1, \dots, d_k)$ and $\underline{e} := (e_1, \dots, e_{k+1})$ be a pair of non-decreasing sequences of positive integers that satisfy the conditions (2), (3), (4). For generic choice of degree- e_j homogeneous polynomials g_j and degree- $(d_i - e_j)$ homogeneous polynomials $h_{i,j}$, where $j = 1, \dots, k+1$ and $i = 1, \dots, \alpha(j)$,*

- (I) *the intersections $V(f_1, \dots, f_s)$ are smooth in codimension three for $s \leq k-1$,*
- (II) *the threefold $V(f_1, \dots, f_k)$ has exactly $\mathcal{S}(\underline{d}; \underline{e})$ nodes on the surface S as its only singularities.*
- (III) *If $\mathcal{S}(\underline{d}; \underline{e}) > 0$, then the surface S is not a Cartier divisor on the threefold $V(f_1, \dots, f_k)$.*

Proof. Claim (I) follows directly from Lemma 3.4. In particular, Lemma 3.4 implies that the fourfold $V(f_1, \dots, f_{k-1})$ is smooth and we can apply Lemma 3.5 to show that all singularities of the threefold $V(f_1, \dots, f_k)$ are nodes. In this way part (II) of the theorem is also proven.

Suppose that $\mathcal{S}(\underline{d}; \underline{e}) > 0$. By (I) and (II) the assumptions of Prop. 2.4 are fulfilled. Consequently, the threefold has exactly $\mathcal{S}(\underline{d}; \underline{e})$ nodes on the surface S and the latter cannot be a Cartier divisor. $\square$

REFERENCES

- [1] S. Cynk, *Euler characteristic of a complete intersection*. in Complex and Differential Geometry (Hannover 2009), 99–114, Springer Proceedings in Mathematics **8**, Springer-Verlag, Berlin, Heidelberg 2011.
- [2] I. Cheltsov, *Factoriality of nodal three-dimensional varieties and connectedness of the locus of log canonical singularities*, Mat. Sb. 197 (2006), 387–414.
- [3] I. Cheltsov, *Points in projective spaces and applications*. J. Differential Geom. 81 (2009), 575–599.
- [4] I. Cheltsov, *Factorial threefold hypersurfaces*. J. Algebraic Geom. 19 (2010), 781–791.
- [5] I. Cheltsov, J. Park, *Factorial hypersurfaces in $\mathbb{P}^4$ with nodes*. Geom. Dedicata 121 (2006), 205–219.
- [6] W. Fulton, *Intersection theory*. Springer-Verlag, New York, 1984.
- [7] W. Decker, G.-M. Greuel, G. Pfister, and H. Schönemann, SINGULAR 3-1-2 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2010).
- [8] J. Harris, *Algebraic geometry: A First Course*. GTM 133, Springer (1995).
- [9] R. Hartshorne, *Ample subvarieties of algebraic varieties*, Springer Lecture Notes in Math. **156**, 1970.
- [10] R. Hartshorne, *Algebraic geometry*. Graduate Texts in Mathematics **52**, Springer-Verlag, New York, 1977.
- [11] G. Kapustka, *Primitive contractions of Calabi-Yau threefolds. II*. J. Lond. Math. Soc. (2) 79 (2009), no. 1, 259–271.
- [12] D. Kosta, *Factoriality condition of some nodal threefolds in P^4* . Manuscripta Math. 127 (2008), 151–166.
- [13] D. Kosta, *Factoriality of complete intersections in $\mathbb{P}^5$* . Proc. Steklov Inst. Math. 264 (2009), 102–109.
- [14] M. Mella, *Birational geometry of quartic 3-folds. II. The importance of being Q-factorial*. Math. Ann. 330 (2004), 107–126.
- [15] J. W. Milnor, *Singular points of complex hypersurfaces*. Princeton Univ. Press, Princeton 1968.

INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

INSTITUTE OF MATHEMATICS OF THE POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-956 WARSZAWA,
P.O. BOX 21, POLAND

E-mail address: `slawomir.cynk@uj.edu.pl`

INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, UL. ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

E-mail address: `slawomir.rams@uj.edu.pl`